



## Improving management of white mold in dry beans: Comparative fungicide efficacy: ProPulse

---

**Michael Wunsch**

North Dakota State University Carrington Research Extension Center

# Improving white mold management in dry beans:

## Comparative fungicide efficacy – methods

---

**Market class** = pinto in most studies; kidney in some studies

**Row spacing** = 14 inches

**Seeding rate** = 90,000 viable seeds/ac in most studies; sometimes 80,000 viable seeds/ac

**Fungicide spray volume** = 15 gal/ac.

Fungicides applied with a hand-held boom pressurized by CO<sub>2</sub>.

**Fungicide spray droplet size:** fine or medium in studies conducted from 2010-2021; fine, medium or coarse, calibrated relative to canopy characteristics, from 2022-2024.

**Number of fungicide applications:** two

**Application timing, first fungicide application:** early bloom and initial pin pod-pod

**Interval between fungicide applications:** 7 to 14 days later, depending on study

**Number of experimental replicates** = 5 or 6 replicates (most studies)

**White mold assessment:** Assessed at/ near dry bean maturity by evaluating every plant individually in for percent of the plant impacted by white mold in a minimum half of the plot.

**Harvest:** To ensure that variability in dry bean standability did not bias yields, plants were clipped at base concurrent with disease assessments, wind-rowed to dry, and manually lifted into the combine.

**Supplemental irrigation:** Supplemental overhead irrigation was applied as needed to establish the white mold disease pressure needed to evaluate fungicide performance.

# **1. Fungicides and fungicide application rates that can be applied twice in-season**

Testing was conducted with two sequential applications of the same fungicide with the goal of rigorously assessing comparative efficacy.

These comparative efficacy results are provided to help facilitate informed decisions for selecting products for application once or twice in-season, either alone or in rotation with another fungicide.

# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 11-13 days later

**ProPulse 10.3 fl oz vs. 8.6 fl oz**

**Combined analysis across eight studies**

Carrington and Langdon, ND (2010, 2012, 2013, 2014)

	<b>WHITE MOLD</b> Severity index % of canopy	<b>DRY BEAN</b> <b>YIELD</b> lbs/ac
Non-treated control	<b>57</b> b	<b>2367</b> b
<b>ProPulse 8.6 fl oz/ac</b>	<b>42</b> a	<b>2816</b> a
<b>ProPulse 10.3 fl oz/ac</b>	<b>41</b> a	<b>2867</b> a
CV:	6.9	5.6

*Within-column means followed by different letters are significantly different. ( $P < 0.05$ ; Tukey procedure).*



# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 12 or 13 days later

## Topsin 30 fl oz/ac vs. ProPulse 10.3 fl oz/ac

Combined analysis across seven studies

Carrington and Langdon, ND (2012, 2014, 2020, 2022)

### WHITE MOLD

Severity index

% of canopy

### DRY BEAN

YIELD

lbs/ac

Non-treated control

**59** b

**2182** b

**Topsin** 30 fl oz/ac

**42** a

**2699** a

**ProPulse** 10.3 fl oz/ac

**44** a

**2754** a

CV:

9.7

5.8

*Within-column means followed by different letters are significantly different. ( $P < 0.05$ ; Tukey procedure).*

# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 12 or 13 days later

## Topsin 30 fl oz/ac vs. ProPulse 8.6 fl oz/ac

Combined analysis across seven studies

Carrington and Langdon, ND (2012, 2014)

### WHITE MOLD

Severity index

% of canopy

### DRY BEAN

YIELD

lbs/ac

Non-treated control

**53** b

**2314** b

**Topsin** 30 fl oz/ac

**34** a

**2789** a

**ProPulse** 8.6 fl oz/ac

**37** a

**2718** a

CV:

12.3

6.9

*Within-column means followed by different letters are significantly different. ( $P < 0.05$ ; Tukey procedure).*

# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 6-14 days later

## Endura 8 oz/ac vs. ProPulse 10.3 fl oz/ac

Combined analysis across 20 studies

Carrington and Langdon, ND (2012, 2013, 2014, 2017, 2020, 2021, 2022, 2023, 2024)

	WHITE MOLD Severity index % of canopy	DRY BEAN YIELD lbs/ac
Non-treated control	<b>49</b> b	<b>2166</b> b
<b>Endura 8.0 oz/ac</b>	<b>35</b> a	<b>2764</b> a
<b>ProPulse 10.3 fl oz/ac</b>	<b>36</b> a	<b>2689</b> a
CV:	14.2	7.4

Within-column means followed by different letters are significantly different. ( $P < 0.05$ ; Tukey procedure).

# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 11-13 days later

## Endura 8 oz/ac vs. ProPulse 8.6 fl oz/ac

### Combined analysis across ten studies

Carrington and Langdon, ND (2012, 2013, 2014, 2023)

	WHITE MOLD Severity index % of canopy	DRY BEAN YIELD lbs/ac
Non-treated control	<b>52</b> b	<b>2300</b> b
<b>Endura 8.0 oz/ac</b>	<b>32</b> a	<b>2924</b> a
<b>ProPulse 8.6 fl oz/ac</b>	<b>39</b> a	<b>2741</b> a
CV:	16.8	7.6

*Within-column means followed by different letters are significantly different. ( $P < 0.05$ ; Tukey procedure).*

# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 12-14 days later

## Omega 13.6 fl oz vs. ProPulse 10.3 fl oz

### Combined analysis across seven studies

Carrington and Langdon, ND (2012, 2013, 2014)

	<b>WHITE MOLD</b> Severity index % of canopy	<b>DRY BEAN</b> <b>YIELD</b> lbs/ac
Non-treated control	<b>50</b> b	<b>2267</b> b
<b>ProPulse</b> 10.3 fl oz/ac	<b>31</b> a	<b>2871</b> a
<b>Omega</b> 13.6 fl oz/ac	<b>24</b> a	<b>3078</b> a
CV:	23.8	10.6

Within-column means followed by different letters are significantly different. ( $P < 0.05$ ; Tukey procedure).



# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 12-13 days later

## Omega 13.6 fl oz vs. ProPulse 8.6 fl oz

Combined analysis across eight studies

Carrington and Langdon, ND (2012, 2014, 2023)

	WHITE MOLD Severity index % of canopy	DRY BEAN YIELD lbs/ac
Non-treated control	<b>49</b> c	<b>2344</b> b
<b>ProPulse</b> 8.6 fl oz/ac	<b>37</b> b	<b>2782</b> a
<b>Omega</b> 13.6 fl oz/ac	<b>24</b> a	<b>3071</b> a
CV	21.7	10.8

Within-column means followed by different letters are significantly different. ( $P < 0.05$ ; Tukey procedure).

# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 12 or 13 days later

## Topsin 30 fl oz/ac vs. Omega 8 fl oz/ac

Combined analysis across three studies

Carrington and Langdon, ND (2014)

*Within-column means followed by different letters  
are significantly different. ( $P < 0.05$ ).*

	WHITE MOLD Severity index % of canopy	DRY BEAN YIELD lbs/ac
Non-treated control	<b>63</b> c	<b>1813</b> c
<b>ProPulse</b> 10.3 fl oz/ac	<b>46</b> b	<b>2373</b> b
<b>Topsin</b> 30 fl oz/ac	<b>40</b> ab	<b>2508</b> ab
<b>Omega</b> 8 fl oz/ac	<b>37</b> ab	<b>2510</b> ab
<b>Endura</b> 8 oz/ac	<b>37</b> ab	<b>2584</b> ab
<b>Omega</b> 13.6 fl oz/ac	<b>25</b> a	<b>2914</b> a

CV:

13.2

6.5

# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 12 or 13 days later

**ProPulse 10.3 fl oz vs. ProPulse 10.3 fl oz + Badge 2 pt**

**Combined analysis across two studies**

Carrington, ND (2020, 2022)

	<b>WHITE MOLD</b> Severity index % of canopy	<b>DRY BEAN</b> <b>YIELD</b> lbs/ac
Non-treated control	<b>82</b> b	<b>1670</b> b
<b>ProPulse 10.3 fl oz/ac</b>	<b>76</b> ab	<b>2137</b> ab
<b>ProPulse 10.3 fl oz + Badge 2 pt/ac</b>	<b>67</b> a	<b>2453</b> a
CV:	2.7	6.1

*Within-column means followed by different letters are significantly different. ( $P < 0.05$ ; Tukey procedure).*

# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 7-14 days later

## Miravis Neo 13.7 fl oz vs. ProPulse 10.3 fl oz

Combined analysis across four studies

Carrington, ND (2022, 2024)

	WHITE MOLD Severity index % of canopy	DRY BEAN YIELD lbs/ac
Non-treated control	<b>70</b> b	<b>1776</b> b
Endura 8.0 oz/ac	<b>58</b> a	<b>2466</b> a
ProPulse 10.3 fl oz/ac	<b>57</b> a	<b>2402</b> a
Miravis Neo 13.7 fl oz/ac	<b>69</b> b	<b>1901</b> b
CV: 10.4		9.3

Within-column means followed by different letters are significantly different. ( $P < 0.05$ ; Tukey procedure).

# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 6-14 days later

## Proline 5.7 fl oz vs. ProPulse 10.3 fl oz

Combined analysis across eight studies

Carrington and Langdon, ND (2009, 2010, 2012, 2013, 2014)

	WHITE MOLD Severity index % of canopy	DRY BEAN YIELD lbs/ac
Non-treated control	<b>58</b> c	<b>2305</b> b
<b>ProPulse</b> 10.3 fl oz/ac	<b>39</b> a	<b>2930</b> a
<b>Proline</b> 5.7 fl oz/ac	<b>48</b> b	<b>2723</b> a
CV:	8.6	8.0

*Within-column means followed by different letters are significantly different. ( $P < 0.05$ ; Tukey procedure).*



# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 6-14 days later

## Proline 5.7 fl oz vs. ProPulse 8.6 fl oz

Combined analysis across six studies

Carrington and Langdon, ND (2010, 2012, 2013, 2014)

	WHITE MOLD Severity index % of canopy	DRY BEAN YIELD lbs/ac
Non-treated control	<b>52</b> b	<b>2628</b> a
<b>ProPulse</b> 8.6 fl oz/ac	<b>39</b> a	<b>2939</b> a
<b>Proline</b> 5.7 fl oz/ac	<b>47</b> b	<b>2895</b> a
CV:	8.6	8.0

Within-column means followed by different letters are significantly different. ( $P < 0.05$ ; Tukey procedure).

# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 13-14 days later

## Quash 2.5 oz vs. ProPulse 8.6 fl oz

Combined analysis across four studies

Langdon and Carrington, ND (2012, 2013)

	WHITE MOLD Severity index % of canopy	DRY BEAN YIELD lbs/ac
Non-treated control	<b>40</b> b	<b>2608</b> b
Endura 8.0 oz/ac	<b>23</b> a	<b>3178</b> a
ProPulse 10.3 fl oz/ac	<b>21</b> a	<b>3244</b> a
Quash 2.5 oz/ac	<b>36</b> b	<b>2751</b> b
CV:	13.8	5.3

Within-column means followed by different letters are significantly different. ( $P < 0.05$ ; Tukey procedure).

# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of the same fungicide,  
initial pin-pod + 13-14 days later

**Rovral 2 pt vs. Switch 14 oz vs. ProPulse 10.3 fl oz**

Combined analysis across four studies

Carrington and Langdon, ND (2012, 2013)

	WHITE MOLD Severity index % of canopy	DRY BEAN YIELD lbs/ac
Non-treated control	<b>40</b> b	<b>2608</b> b
Endura 8.0 oz/ac	<b>23</b> a	<b>3178</b> a
ProPulse 10.3 fl oz/ac	<b>21</b> a	<b>3244</b> a
Rovral 2.0 pt/ac	<b>27</b> a	<b>3015</b> a
Switch 14 oz/ac	<b>26</b> a	<b>2979</b> a
CV:	17.7	4.4

Within-column means followed by different letters are significantly different. ( $P < 0.05$ ; Tukey procedure).

## **2. Fungicides and fungicide application rates that can only be applied once in-season**

Testing was conducted with two sequential applications of Topsin/generic at 40 fl oz/ac in order to fit the testing within existing protocols in which products were applied twice sequentially.

Comparative efficacy results are provided for Topsin/generic at 40 fl oz/ac to help facilitate informed decisions for selecting products **for application once in-season, either as a single application or in rotation with another fungicide.**

# Comparative fungicide efficacy: white mold in dry edible beans

## Topsin, 40 fl oz/ac vs. ProPulse, 10.3 fl oz/ac

### Combined analysis across four studies

Carrington and Langdon, ND (2012, 2023, 2024)

	WHITE MOLD Severity index % of canopy	DRY BEAN YIELD lbs/ac	
Non-treated control	<b>51</b> b	<b>2243</b> b	<i>Within-column means followed by different letters are significantly different</i>  ( $P < 0.05$ ; Tukey procedure).
<b>Topsin</b> 40 fl oz/ac	<b>33</b> a	<b>3052</b> a	
<b>ProPulse</b> 10.3 fl oz/ac	<b>36</b> a	<b>2830</b> a	
CV:	9.4	5.9	

### Read the label for Topsin/generic carefully.

The labels for some brands of the flowable formulation of thiophanate-methyl explicitly state that the product can only be applied once per season at 40 fl oz/ac. When applied twice, the maximum application rate is 30 fl oz/ac.

The labels for other brands of the flowable formulation of thiophanate-methyl lack explicit language prohibiting two applications at 40 fl oz/ac but indicate a usage rate of 30-40 fl oz/ac when applied once and 20-30 fl oz/ac when applied two or more times (max. 80 fl oz/ac). **Two applications at 40 fl oz/ac in the same season should be considered off-label.**

In these studies, same fungicide was applied twice sequentially 6-13 days apart in order to fit the testing within existing protocols in which products were applied twice sequentially. Comparative efficacy data of Topsin/generic applied at 40 fl oz/ac are provided to inform decision-making when applying Topsin/generic once during the season, either as a stand-alone application or in rotation with another fungicide.



# Comparative fungicide efficacy: white mold in dry edible beans

Two sequential applications of same fungicide vs. fungicide rotation,  
initial pin-pod + 7-14 days later

**Topsin 30 fl oz f.b. Endura 8 oz vs.**

**Topsin 30 fl oz rotated with ProPulse 10.3 fl oz**

Combined analysis across six studies

Carrington, ND (2015, 2017, 2019, 2022, 2023)

**WHITE MOLD**  
Severity index  
% of canopy

**DRY BEAN**  
**YIELD**  
lbs/ac

Non-treated control

**50** b

**1904** b

**Topsin 30 fl oz/ac / Endura 8 oz/ac**

**42** a

**2356** a

**Topsin 30 fl oz / ProPulse 10.3 fl oz/ac**

**40** a

**2395** a

**ProPulse 10.3 fl oz / Topsin 30 fl oz/ac**

**40** a

**2470** a

CV:

11.3

7.8

*Within-column means followed by different letters are significantly different ( $P < 0.05$ ; Tukey procedure).*

## Improving white mold management in dry beans:

# Comparative fungicide efficacy versus ProPulse

---

## Conclusions from comparative efficacy testing

### **Most effective fungicides: two applications/season**

Omega at 13.6 fl oz/ac, Endura at 8 oz/ac, ProPulse at 10.3 fl oz/ac, Topsin at 30 fl oz/ac

### **Most effective fungicides: if only a single application is made**

Omega at 13.6 fl oz/ac, Topsin at 40 fl oz/ac, Endura at 8 oz/ac, ProPulse at 10.3 fl oz/ac, Topsin at 30 fl oz/ac

### **Optimal fungicide rotation sequence with Topsin/generic**

ProPulse applied first, Topsin applied second

### **Less effective fungicides**

Proline at 5.7 fl oz, Quash at 2.5 oz/ac, Miravis Neo 13.6 fl oz/ac, Rovral 2 pt/ac, Switch 14 oz/ac



## People

Staff, Carrington: Aaron Fauss, Suanne Kallis, Jesse Hafner, Gabriela Henson, Thomas Miorini, Billy Kraft, Michael Schaefer

Staff in Langdon who contributed to this research: Scott Halley, Amanda Arens





## Funding support

### Research funding:

- Northarvest Bean Growers Association
- ND Crop Protection Product Harmonization & Registration Board
- Contract testing (BASF, Bayer, Corteva, FMC, Gowan, Syngenta, Valent, others)

### Seed was donated by:

- Bollingberg Seeds Company (Kurt Bollingburg; Cathay, ND)
- Green Valley Bean Company (John Berthold; Park Rapids, MN)
- Kelley Bean Company; Hatton, ND (Dean Nelson)