

# Improving management of white mold in soybeans: Comparative fungicide efficacy – ProPulse

Improving white mold management in soybeans:

Comparative fungicide efficacy – fundamental concepts

# This is an abbreviated summary of comparative fungicide efficacy in field trials conducted from 2019-2024.

Included are all studies conducted 2019 to 2024. Starting in 2019, fungicide spray droplet size was calibrated relative to soybean canopy closure in all soybean white mold fungicide efficacy testing in Carrington and Oakes, ND.

Full results are available in the accompanying PDF fungicide efficacy testing results posted online at <u>https://www.ndsu.edu/agriculture/ag-hub/research-extension-centers-recs/carrington-rec/research/plant-pathology</u> (or search for 'NDSU Carrington plant pathology').

Improving white mold management in soybeans: Comparative fungicide efficacy – fundamental concepts

# Calibrating fungicide droplet size relative to soybean canopy closure is critical.

Calibrating fungicide droplet size relative to canopy closure sharply improves the performance of all fungicides for white mold management in soybeans and is particularly important for fungicides with intermediate efficacy.

The magnitude of the difference between the most effective fungicides and fungicides with intermediate efficacy is reduced when spray droplet size is optimized relative to soybean canopy closure.

Soybean row spacing = 14".

Seeding rate = 140,000 or 165,000 viable seeds/ac.

**Spray volume** = 15 gal/ac.

Fungicides applied with a hand-held boom pressurized by  $CO_2$ .

**Spray droplet size:** Fungicides applied with TeeJet nozzles emitting fine droplets when average canopy closure < 80%, medium droplets when average canopy closure was between 80 and 90%, medium-coarse or coarse droplets when average canopy closure was between 90 and 97%, and coarse droplets when canopy closure average 97-100%.

#### **Number of experimental replicates** = 6

**White mold assessment:** Assessed at soybean maturity by evaluating every plant individually in 1 or 2 rows/plot (entire length of row) for the percent of the plant impacted by white mold.

**Application timing:** when conditions favor white mold as soybeans enter bloom, 100% R2 or canopy closure, whichever occurred first; when conditions did not favor white mold as soybeans entered bloom, early/full R3; second application (when made) 7 to 14 days later.

**Supplemental irrigation:** Supplemental overhead irrigation was applied as needed to establish the white mold disease pressure needed to evaluate fungicide performance.

#### **TWO FUNGICIDE APPLICATIONS**

# **Delaro Complete** 8 fl oz/ac > **ProPulse** 6 fl oz/ac

#### COMBINED ANALYSIS ACROSS 3 STUDIES

Carrington, ND (2022, 2023, 2024)

ProPulse: fluopyram (FRAC 7) + prothioconazole (FRAC 3)

#### Two fungicide applications 7 or 9 or 11 days apart

Delaro Complete: fluopyram (FRAC 7), prothioconazole (FRAC 3), trifloxystrobin (FRAC 11)

|                            | White mold  | White mold     | Soybean     |
|----------------------------|-------------|----------------|-------------|
|                            | Incidence   | Severity Index | Yield       |
|                            | % plants    | % canopy       | bu/ac       |
| Non-treated control        | <b>61</b> b | <b>40</b> b    | <b>43</b> a |
| ProPulse 6 fl oz/ac        | <b>37</b> a | <b>22</b> a    | <b>51</b> a |
| Delaro Complete 8 fl oz/ac | <b>34</b> a | <b>20</b> a    | <b>53</b> a |
| C/                         | V: 15.3     | 4.4            | 10.9        |

#### **Michael Wunsch**

#### **TWO FUNGICIDE APPLICATIONS**

### Endura 6 oz/ac > ProPulse 6 fl oz/ac

| COMBINED | ANALYSIS, | 6 STUDIES |
|----------|-----------|-----------|
|----------|-----------|-----------|

Carrington, ND (2022, 2023, 2024)

Endura: boscalid (FRAC 7)

#### Two sequential fungicide applications 7 or 9 or 11 days apart

Delaro Complete: fluopyram (FRAC 7), prothioconazole (FRAC 3), trifloxystrobin (FRAC 11)

|                     | White mold  | White mold     | Soybean     |
|---------------------|-------------|----------------|-------------|
|                     | Incidence   | Severity Index | Yield       |
|                     | % plants    | % canopy       | bu/ac       |
| Non-treated control | <b>74</b> b | <b>57</b> b    | <b>33</b> b |
| Endura 6 oz/ac      | <b>47</b> a | <b>33</b> a    | <b>45</b> a |
| ProPulse 6 fl oz/ac | <b>52</b> a | <b>38</b> a    | <b>44</b> a |
| CV                  | 2: 8.4      | 13.9           | 10.3        |

#### **Michael Wunsch**

#### **TWO FUNGICIDE APPLICATIONS**

# Viatude 16 fl oz/ac > ProPulse 6 fl oz/ac

| COMBINED ANALYSIS, 3 STUD                                      | IES                   |                |  |  |
|--|-----------------------|----------------|--|--|
| Carrington, ND (2024)  |                       |                | Two sequential fungicide applications 7 days apart |  |
| <b>ProPulse:</b> fluopyram (FRAC 7) + prothioconazole (FRAC 3) |                       |                |  |  |
| Viatude: picoxystrobin (FRAC 11)                               | + prothioconazole (FR | RAC 3)         |  |  |
|  | White mold            | White mold     | Soybean  |  |
|  | Incidence             | Severity Index | Yield  |  |
|  | % plants              | % canopy       | bu/ac  |  |
| Non-treated control  | <b>87</b> b           | <b>74</b> b    | <b>24</b> b  |  |
| ProPulse 6 fl oz/ac  | <b>68</b> a           | <b>53</b> a    | <b>37</b> a  |  |
| Viatude 16 fl oz/ac  | <b>62</b> a           | <b>47</b> a    | <b>39</b> a  |  |
| CV   | /: 7.5                | 8.5            | 4.2  |  |

#### **Michael Wunsch**

Improving white mold management in soybeans:

Comparative fungicide efficacy – ProPulse

#### SINGLE FUNGICIDE APPLICATION

# ProPulse 8 fl oz > Delaro Complete 8 fl oz > ProPulse 6 fl oz

| COMBINED ANALYSIS ACROSS 4 STUDIES<br>Carrington, ND (2020, 2022, 2023 2024)             | Single fungicide application |  |  |
|--|------------------------------|--|--|
| Delaro Complete: fluopyram (FRAC 7), trifloxystrobin (FRAC 11), prothioconazole (FRAC 3) |                              |  |  |
| ProPulse: fluopyram (FRAC 7), prothioconazole (FRAC 3)                                   |                              |  |  |

|                            | White mold   | White mold     | Soybean      |
|----------------------------|--------------|----------------|--------------|
|                            | Incidence    | Severity Index | Yield        |
|                            | % plants     | % canopy       | bu/ac        |
| Non-treated control        | <b>66</b> b  | <b>50</b> b    | <b>37</b> b  |
| Delaro Complete 8 fl oz/ac | <b>53</b> ab | <b>34</b> a    | <b>45</b> a  |
| ProPulse 6 fl oz/ac        | <b>59</b> ab | <b>40</b> ab   | <b>43</b> ab |
| ProPulse 8 fl oz/ac        | <b>49</b> a  | <b>31</b> a    | <b>46</b> a  |
| C\                         | /: 10.5      | 18.2           | 7.6          |

#### **Michael Wunsch**

#### SINGLE FUNGICIDE APPLICATION

# ProPulse 8 fl oz/ac > Endura 5.5 oz/ac > ProPulse 6 fl oz/ac

| COMBINED ANALYSIS ACROSS 3 S                                  | TUDIES                  |                              | lala ann Raathan |  |
|---|-------------------------|------------------------------|------------------|--|
| Carrington, ND (2022, 2023 2024)                              |                         | Single fungicide application |                  |  |
| <b>ProPulse:</b> fluopyram (FRAC 7), prothioconazole (FRAC 3) |                         |                              |                  |  |
| Endura: boscalid (FRAC 7)                                     |                         |                              |                  |  |
|   | White mold<br>Incidence | White mold<br>Severity Index | Soybean<br>Yield |  |

|                     | %   | 6 plants     | % canopy     | bu/ac       |
|---------------------|-----|--------------|--------------|-------------|
| Non-treated control |     | <b>75</b> b  | <b>61</b> b  | <b>32</b> b |
| ProPulse 6 fl oz/ac |     | <b>63</b> ab | <b>46</b> ab | <b>40</b> a |
| ProPulse 8 fl oz/ac |     | <b>55</b> a  | <b>39</b> a  | <b>43</b> a |
| Endura 5.5 oz/ac    |     | <b>61</b> ab | <b>43</b> a  | <b>42</b> a |
|                     | CV: | 9.5          | 12.5         | 6.3         |

#### **Michael Wunsch**

# Approximate comparative efficacy ranking among those fungicides assessed relative to ProPulse:

- 1. ProPulse, 8 fl oz/ac
- 2. Delaro Complete, 8 fl oz/ac
- 3. Endura, 6 oz/ac
- 4. Viatude, 16 fl oz/ac
- 5. ProPulse, 6 fl oz/ac

# **Special thanks:**



#### People:

Aaron Fauss, Suanne Kallis, Jesse Hafner, Gabriela Henson; Student workers

This research was funded by pesticide registrants:

BASF, Bayer, Corteva, Gowan, Syngenta